

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 :
G01N 33/68, C07K 14/47, 14/82**A1****(11) International Publication Number: WO 96/33416****(43) International Publication Date: 24 October 1996 (24.10.96)****(21) International Application Number: PCT/US96/05639****(22) International Filing Date: 19 April 1996 (19.04.96)****(30) Priority Data:**
08/426,529 20 April 1995 (20.04.95) US**(60) Parent Application or Grant**
(63) Related by Continuation
US 08/426,529 (CIP)
Filed on 20 April 1995 (20.04.95)**(71) Applicant (for all designated States except US):** LXR
BIOTECHNOLOGY INC. [US/US]; 1401 Marina Way
South, Richmond, CA 94804-3746 (US).**(72) Inventors; and****(75) Inventors/Applicants (for US only):** BARR, Philip, J. [US/US];
152 Hillcrest Road, Berkley, CA 94705 (US). KIEFER,
Michael, C. [US/US]; 401 Wright Court, Clayton, CA 94517
(US).**(74) Agents:** LEHNHARDT, Susan, K. et al.; Morrison & Foerster
L.L.P., 755 Page Mill Road, Palo Alto, CA 94303-1018
(US).**(81) Designated States:** AL, AM, AT, AU, AZ, BB, BG, BR, BY,
CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP,
KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MG, MK,
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG,
SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO
patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, ML, MR, NE, SN, TD, TG).**Published***With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.***(54) Title: METHODS OF SCREENING FOR THERAPEUTIC AGENTS USING NOVEL APOPTOSIS-MODULATING PROTEINS****(57) Abstract**

The present invention provides methods to screen for anti-viral agents utilizing a novel family of apoptosis-modulating proteins.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**METHODS OF SCREENING FOR THERAPEUTIC AGENTS USING
NOVEL APOPTOSIS-MODULATING PROTEINS**

TECHNICAL FIELD

5 This invention relates to methods of screening for therapeutic agents using novel proteins with apoptosis-modulating activity.

BACKGROUND ART

10 Apoptosis is a normal physiologic process that leads to individual cell death. This process of programmed cell death is involved in a variety of normal and pathogenic biological events and can be induced by a number of unrelated stimuli. Changes in the biological regulation of apoptosis also occur during aging and are responsible for many of the conditions and diseases related to aging. Recent studies of apoptosis have implied that a common metabolic pathway leading to cell death may be initiated by a wide variety of signals, including
15 hormones, serum growth factor deprivation, chemotherapeutic agents, ionizing radiation and infection by human immunodeficiency virus (HIV). Wyllie (1980) *Nature* 284:555-556; Kanter et al. (1984) *Biochem. Biophys. Res. Commun.* 118:392-399; Duke and Cohen (1986) *Lymphokine Res.* 5:289-299; Tomei et al. (1988) *Biochem. Biophys. Res. Commun.* 155:324-331; Kruman et al. (1991) *J. Cell. Physiol.* 148:267-273; Ameisen and Capron (1991) *Immunology Today* 12:102; and Sheppard and Ascher (1992) *J. AIDS* 5:143. Agents that modulate
20 the biological control of apoptosis thus have therapeutic utility in a wide variety of conditions.

25 Apoptotic cell death is characterized by cellular shrinkage, chromatin condensation, cytoplasmic blebbing, increased membrane permeability and interchromosomal DNA cleavage. Kerr et al. (1992) *FASEB J.* 6:2450; and Cohen and Duke (1992) *Ann. Rev. Immunol.* 10:267. The blebs, small, membrane-encapsulated spheres that pinch off of the surface of apoptotic cells,

may continue to produce superoxide radicals which damage surrounding cell tissue and may be involved in inflammatory processes.

5 The Bcl-2 gene was discovered at the common chromosomal translocation site t(14:18) in follicular lymphomas and results in aberrant over-expression of bcl-2. Tsujimoto et al. (1984) *Science* 226:1097-1099; and Cleary et al. (1986) *Cell* 47:19-28. The normal function of bcl-2 is the prevention of apoptosis; unregulated expression of bcl-2 in B cells is thought to lead to increased numbers of proliferating B cells which may be a critical factor in the development of lymphoma. McDonnell and Korsmeyer (1991) *Nature* 349:254-256; and, for
10 review see, Edgington (1993) *Bio/Tech.* 11:787-792. Bcl-2 is also capable of blocking of γ irradiation-induced cell death. Sentman et al. (1991) *Cell* 67:879-888; and Strassen (1991) *Cell* 67:889-899. It is now known that bcl-2 inhibits most types of apoptotic cell death and is thought to function by regulating an antioxidant pathway at sites of free radical generation. Hockenbery et al. (1993)
15 *Cell* 75:241-251.

Apoptosis, a normal cellular event, can also be induced by pathological conditions and a variety of injuries. Apoptosis is involved in a wide variety of conditions including, but not limited to: cardiovascular disease; cancer regression; immunoregulation; viral diseases; anemia; neurological disorders; gastrointestinal
20 disorders such as diarrhea and dysentery; diabetes; hair loss; rejection of organ transplants; prostate hypertrophy; obesity; ocular disorders; stress; and aging.

Bcl-2 belongs to a family of proteins of which some have been cloned and sequenced. Williams and Smith (1993) *Cell* 74:777-779. Various Bcl-2 members have the ability to associate with one another as heterodimers. Oltvai et al. (1993) *Cell* 74:609-619; and Sato et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:9238-9242.
25 Additionally, BHRF1 displays a 25% sequence identity to Bcl-2 (Cleary et al. (1986) *Cell* 47:19-28) and has been shown by gene transfer studies to protect B cells from apoptosis. Henderson et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:8479-8483.

The herpesvirus family of viruses typically produce latent and recurrent infections. Herpesvirus genomes are composed of sequences with a short and a long region. Herpesvirus particles have a diameter from 180 nm to 200 nm. Many particles do not contain envelopes. Typically, the DNA is wrapped around an associated protein. The herpesvirus has a tendency to persist in a quiescent state for irregular periods of time.

All references cited herein, both *supra* and *infra*, are hereby incorporated by reference herein.

SUMMARY OF THE INVENTION

Methods of screening for pharmaceutical agents that stimulate, as well as pharmaceutical agents that inhibit Bak and Bak-2 protein activity levels are provided. The methods include combining a Bak protein and a viral protein under conditions in which they interact to form a test sample, exposing the test sample to a potential therapeutic agent and monitoring the interaction of the proteins. Potential therapeutic agents which disrupt the interaction compared to control test samples to which no agent has been added are selected for further study.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the Bak cDNA nucleotide sequence and amino acid sequence encoded thereby.

Figure 2 shows the sequence of the Bak-2 cDNA and flanking sequences and the corresponding predicted amino acid sequence of the Bak-2 protein.

Figure 3 shows the interactions of Bak and Flag-Bak (F-Bak) fusion proteins with Epstein-Barr virus BHRF-1 protein. In column A, lane 1 depicts the results obtained from *in vitro* co-translated proteins F-Bak/BHRF-1 and lane 2 depicts F-Bak/BHRF-1 proteins bound to anti-FLAG agarose. In column B, the lanes are the same with the exception that the Bak protein is Bak-2.

DISCLOSURE OF THE INVENTION

The present invention provides methods of screening for potential anti-viral therapeutic agents. The proteins encoded by nucleotide sequences encoding the novel bcl-2 homologs, Bak and Bak-2 proteins have been found to interact with the Epstein Barr Virus (EBV) protein BHRF1 indicating that Bak proteins contribute to the pathogenicity of the disease. BHRF1 is an EBV early lytic cycle protein. Pearson et al. (1987) *Virology* 160:151-161. The invention encompasses methods containing the steps of exposing the Bak proteins and viral proteins, or functional portions thereof, to potential therapeutic agents and monitoring the interaction of the proteins. The invention further utilizes recombinant cells and transgenic animals expressing the cloned Bak or Bak-2 genes.

The cloning and analysis of Bak genes and proteins are described in detail in commonly owned WO application PCT/US94/13930. Bak genes and proteins are also described in Kiefer et al. (1995) *Nature* 374:736. The nucleotide and predicted amino acid residue sequences of Bak protein are shown in Figure 1; and those of Bak-2 are shown in Figure 2. Bak mRNA has been detected in a variety of human organs and tissues by Northern blot analysis. These organs include liver; heart; skeletal muscle; lung; kidney; and pancreas.

These references also disclose that the Bak proteins are capable of modulating apoptosis. In a lymphoblastoid cell line, expression of Bak protein was shown to decrease Fas-mediated apoptosis. In a mouse progenitor B cell line, FL5.12, Bak-2 protein and a derivative of Bak protein decrease IL-3-induced apoptosis whereas Bak protein increased apoptosis. Thus, depending on the cell type, the derivative of Bak protein, and the method of induction of apoptosis, apoptosis can be modulated in a highly specific manner by controlling the concentration of Bak proteins.

As used herein, the term "Bak gene(s)" refers to the nucleic acid molecules described herein and in PCT/US94/13930, "the Bak protein(s)" refers to the proteins encoded thereby. The nucleotides include, but are not limited to, the

cDNA and complementary DNA, genome-derived DNA and synthetic or semi-synthetic DNA or RNA. The nucleotide sequence of the Bak cDNA with the location of restriction endonuclease sites is shown in Figure 1.

5 The nucleotide sequence of Bak-2 cDNA, along with the predicted amino acid sequence of Bak-2 protein and the locations of restriction endonuclease recognition sites, is given in Figure 2. The Bak gene is on human chromosome 6 and the Bak-2 gene is on human chromosome 20. There is also a member of the family, Bak-3, which is on human chromosome 11. Bak-3 appears to be a
10 pseudogene. Fluorescence *in situ* hybridization (FISH) indicated an approximate location of the Bak gene to be at 6p21-23.

The invention includes the use of modified Bak DNA sequences such as deletions, substitutions and additions particularly in the non-coding regions of genomic DNA. Such changes are useful to facilitate cloning and modify gene
15 expression. Any DNA which encodes a portion of a Bak protein sufficient to bind to BHRF1 or any other suitable viral protein is suitable for use herein. As described below, various fusion proteins are suitable for use herein.

Various substitutions can be made within the coding region that either do not alter the amino acid residues encoded or result in conservatively substituted amino acid residues. Nucleotide substitutions that do not alter the amino acid
20 residues encoded are useful for optimizing gene expression in different systems. Suitable substitutions are known to those of skill in the art and are made, for instance, to reflect preferred codon usage in the particular expression systems.

The invention encompasses the use of functionally equivalent variants and derivatives of Bak genes which may enhance, decrease or not significantly affect
25 the properties of Bak proteins. For instance, changes in the DNA sequence that do not change the encoded amino acid sequence, as well as those that result in conservative substitutions of amino acid residues, one or a few amino acid deletions or additions, and substitution of amino acid residues by amino acid analogs are those which will not significantly affect its properties.

Amino acid residues which can be conservatively substituted for one another include but are not limited to: glycine/alanine; valine/isoleucine/leucine; asparagine/glutamine; aspartic acid/glutamic acid; serine/threonine; lysine/arginine; and phenylalanine/tyrosine. Any conservative amino acid substitution which does not significantly affect the properties of Bak proteins is encompassed by the present invention.

Techniques for nucleic acid manipulation useful for the practice of the present invention are described in a variety of references, including, but not limited to, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Vol. 1-3, eds. Sambrook *et al.*, Cold Spring Harbor Laboratory Press (1989); and *Current Protocols in Molecular Biology*, eds. Ausubel *et al.*, Greene Publishing and Wiley-Interscience: New York (1987) and periodic updates.

The coding regions of Bak genes can also be ligated into expression vectors capable of stably integrating into other cell types including but not limited to cardiomyocytes, neural cell lines such as GTI-7 and TNF sensitive cells such as the human colon adenocarcinoma cell line HT29 so as to provide a variety of assay systems to monitor the regulation of apoptosis by Bak proteins.

As used herein, "BHRF1" or "viral proteins" encompasses the full length EBV protein and portions or derivations thereof sufficient to bind to Bak proteins or portions or derivatives thereof. Such proteins include, but are not limited to, homologous proteins expressed by any virus, particularly various forms of herpes and herpes-like viruses, such as cytomegalovirus and varicella zoster.

The interaction between a Bak protein and viral protein such as BHRF-1 can be produced by adding purified proteins together. Preferably, however, the proteins are cotranscribed and translated under conditions that allow protein-protein interactions. Co-translation can be performed in vitro or in vivo in whole cells expressing native or recombinant Bak proteins and viral proteins. Any suitable recombinant expression vectors may be used. The Bak proteins can also

be separately translated and then combined under conditions that allow for protein-protein interactions.

5 Methods of monitoring protein interactions are known in the art, any method is suitable for use herein. Preferably, co-precipitation is used. The ability of an antibody to precipitate one of the proteins or an immunological tag fused thereto is used to immunoprecipitate the protein and the immunoprecipitate is monitored for the presence of both proteins. Methods of co-precipitation are known in the art and are described in the examples below. Any other method in the art is suitable for use herein, including, but not limited to, protein interactive trapping, such as GST fusion protein immobilization on glutathione columns and, ELISA. Immunological tags are often incorporated into fusion proteins and including, for instance, FLAG, hemagglutinin and glutathione-S transferase.

10 Purification or isolation of Bak proteins expressed either by the recombinant DNA or from biological sources such as tissues can be accomplished by any method known in the art. Protein purification methods are known in the art. Generally, substantially purified proteins are those which are free of other, contaminating cellular substances, particularly proteins. Preferably, the purified Bak proteins are more than eighty percent pure and, most preferably, more than ninety-five percent pure. For clinical use as described below, the Bak proteins are preferably highly purified, at least about ninety-nine percent pure, and free of pyrogens and other contaminants.

20 Suitable methods of protein purification are known in the art and include, but are not limited to, affinity chromatography, immunoaffinity chromatography, size exclusion chromatography, HPLC and FPLC. Any purification scheme that does not result in substantial degradation of the protein is suitable for use herein.

25 As used herein, "Bak proteins" includes functionally equivalent variants thereof which do not significantly affect their properties and variants which retain the same overall amino acid sequence but which have enhanced or decreased activity. For instance, conservative substitutions of amino acid residues, one or a

few amino acid deletions or additions, and substitution of amino acid residues by amino acid analogs are within the scope of the invention.

Amino acid residues which can be conservatively substituted for one another include but are not limited to: glycine/alanine; valine/isoleucine/leucine; 5 asparagine/glutamine; aspartic acid/glutamic acid; serine/threonine; lysine/arginine; and phenylalanine/tyrosine. Any conservative amino acid substitution which does not significantly affect the properties of Bak proteins is encompassed by the present invention.

Suitable antibodies for use herein are generated by using the Bak proteins 10 as an antigen or, preferably, peptides encompassing the Bak protein regions that lack substantial homology to the other gene products of the bcl family. Antibodies to the viral proteins are also suitable for use herein. Methods of detecting proteins using antibodies and of generating antibodies using proteins or synthetic peptides are known in the art and are not described in detail herein.

15 Screening for therapeutically effective agents is done by exposing the Bak protein and the viral protein to such agents which may directly or indirectly affect the interaction between a Bak protein and a viral protein. Suitable potential therapeutic agents include, but are not limited to, any pharmaceutical agent such as cytokines, small molecule drugs, cell-permeable small molecule drugs, 20 hormones, combinations of interleukins, lectins and other stimulating agents, e.g., PMA, LPS, bispecific antibodies, peptide mimetics, antisense oligonucleotides and other agents which modify cellular functions or protein expression.

The proteins are added together or co-expressed, exposed to such agents at physiologically effective concentrations, and the interaction thereof is measured 25 relative to a control not exposed to such agents. Those biological modifiers which decrease the interaction between a Bak protein and a viral protein relative to a control are selected for further study of their anti-viral activity.

As previously shown, overexpressed Bak proteins protect EBV-transformed B cells from apoptosis following serum withdrawal or anti-Fas

5 treatment. PCT/US94/13930. These results indicate that a Bak-BHRF1 interaction exists whereby BHRF1 not only neutralizes the normally apoptotic effect of Bak protein, but additionally induces a protective activity. Alternatively, propagation of cells transfected with the Bak cDNA might select for cells that are expressing high levels of BHRF1 or other EBV encoded anti-apoptotic proteins. This could lead to an anti-apoptotic response upon subjecting the cells to an apoptosis signal such as serum withdrawal. Example 2 shows that *in vitro* translated Flag-Bak (epitope tagged) and BHRF1 can be coprecipitated with an antibody that recognizes the Flag epitope indicating that Bak proteins and BHRF1
10 interact directly with one another.

The following examples are provided to illustrate but not limit the present invention. Unless otherwise specified, all cloning techniques were essentially as described by Sambrook et al. (1989) and all reagents were used according to the manufacturer's instructions.

15 Example 1

Expression of Recombinant Bak Gene

In order to express the recombinant Bak gene in the baculovirus system, the Bak cDNA generated as described in PCT/US94/13930 was used to generate a novel Bak vector, by PCR, using primers from the 3' and 5' flanking regions of the
20 gene which contain restriction sites to facilitate cloning. The plasmids were sequenced by the dideoxy terminator method (Sanger *et al.*, 1977) using sequencing kits (USB, Sequenase version 2.0) and internal primers. This was to confirm that no mutations resulted from PCR.

25 A clone was used to generate recombinant viruses by *in vivo* homologous recombination between the overlapping sequences of the plasmid and AcNPV wild type baculovirus. After 48 hours post-transfection in insect *Spodoptera frugiperda* clone 9 (SF9) cells, the recombinant viruses were collected, identified by PCR and further purified. Standard procedures for selection, screening and propagation of recombinant baculovirus were performed in accordance with the

manufacturer's instructions (Invitrogen). The molecular mass, on sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), of the protein produced in the baculovirus system was compared with the predicted molecular mass of Bak protein according to the amino-acid sequence.

5 In addition, similar clones can be expressed in any expression system known in the art including, but not limited to, bacterial, yeast, insect and mammalian. A suitable yeast intracellular expression system is described by Barr et al. (1992) *Transgenesis* ed. JAH Murray, (Wiley and Sons) pp. 55-79.

10 The Bak gene coding sequence was excised and introduced into plasmids pCEP7, pREP7 and pcDNA3 (Invitrogen) at compatible restriction enzyme sites. pCEP7 was generated by removing the RSV 3'-LTR of pREP7 with *Xba*I/*Asp*718, and substituting the CMV promoter from pCEP4 (Invitrogen). 25 µg of each Bak-containing plasmid was electroporated into the B lymphoblastoid cell line WIL-2, and stable hygromycin resistant transformants or G418 resistant
15 transformants (pcDNA3 constructs) expressing Bak were selected.

Example 2

Bak proteins interact with Epstein-Barr Virus encoded BHRF1 protein

BHRF1 cDNA was amplified by RT-PCR from WI-L2 mRNA using standard PCR protocol according to the instructions of the manufacturers of the
20 PCR kit and thermal cycler (Perkin Elmer Cetus). The Flag-Bak and Flag-Bak-2 cDNAs were generated by RT-PCR as above from Bak and Bak-2/pcDNA3 plasmid templates but included the 24 base Flag encoding sequence 5'-GAC TAC AAG GAC GAC GAT GAC AAG-3' in the sense primer. This resulted in a cDNA encoding N-terminal Flag-Bak and Flag-Bak-2 fusion proteins that could
25 be recognized and purified by the anti-Flag M2 antibody (Kodak-IBI). The cDNAs were ligated into the pcDNA3 vector which is under the control of the CMV and T7 RNA polymerase promoter. The Flag-Bak and BHRF1 plasmids or Flag-Bak-2 and BHRF1 plasmids were then cotranscribed and cotranslated using

the TnT coupled reticulocyte lysate system according to manufacturer's instructions (Promega).

Briefly, 0.5-1.0 μ g of the two circular plasmids were simultaneously transcribed and translated in 50 μ L of TnT lysate for 90 minutes at 32°C. After translation, 20 μ L of lysate was mixed with 20 μ L of 2X PBS plus 40 μ L of anti-Flag M2 affinity gel (Kodak) and incubated with gentle rocking overnight at 4°C. Immunoprecipitates were collected by centrifugation in an Eppendorf microfuge at 1500 rpm for 15 minutes at 4°C. Pellets were washed 4 times with 1.5 mL PBS and after the final wash were resuspended in 30 μ L of SDS-PAGE sample buffer. The samples were then analyzed by SDS-PAGE on a 18% polyacrylamide gel. Gels were fixed with 10% glacial acetic acid, dried and exposed to X-ray film overnight at room temperature.

As shown in Figure 3, Flag-Bak and BHRF1 as well as Flag-Bak-2 and BHRF1 were efficiently cotranscribed and cotranslated (lanes 1). Clearly, the anti-Flag M2 antibody effectively coprecipitates Flag-Bak and BHRF1 or Flag-Bak-2 and BHRF1 (lanes 2). This demonstrates that BHRF1 interacts with both Bak and Bak-2 proteins *in vitro* and suggests that such interactions occur *in vivo* resulting in the modulation of apoptosis. Interactions of Bak proteins with viral proteins are likely to have evolved to allow viral replication or latency to proceed in the absence of apoptotic death of the host cell. Interference, therefore, in these interactions represents an important new strategy for the design of novel antiviral agents. Similarly, malignant cells derived from transformation by viruses such as EBV would also be amenable to diagnosis or therapy with these agents.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be apparent to those skilled in the art that certain changes and modifications may be practiced. Therefore, the description and examples should not be construed as limiting the scope of the invention, which is delineated by the appended claims.

CLAIMS

We claim:

- 5 1. A method for screening potential anti-viral therapeutic agents comprising the steps of:
- (a) combining a Bak protein and a viral protein under conditions in which they interact, to form a test sample;
 - (b) exposing the test sample to a potential therapeutic agent and;
 - 10 (c) monitoring the interaction of the Bak protein and the viral protein; wherein, a potential therapeutic agent is selected for further study when it disrupts the interaction compared to a control test sample to which no potential therapeutic agent has been added.
- 15 2. The method according to claim 1, wherein the potential therapeutic agent is selected from the group consisting of any pharmaceutical agent, cytokines, small molecule drugs, cell-permeable small molecule drugs, hormones, combinations of interleukins, lectins and other stimulating agents e.g. PMA, LPS,
- 20 bispecific antibodies, peptide mimetics, antisense oligonucleotides and other agents which modify cellular functions or protein expression.
- 25 3. The method according to claim 1, wherein the Bak protein is selected from the group consisting of Bak, Bak-2, portions thereof sufficient to effect binding to a viral protein and fusion proteins thereof containing a portion thereof sufficient to effect binding to a viral protein.

4. The method according to claim 3, wherein the fusion protein is selected from the group consisting of epitope-tagged Bak protein and epitope-tagged Bak-2 protein.

5 5. The method according to claim 1, wherein the viral protein is selected from the group consisting of Epstein Barr Virus BHRF1 protein, other EBV encoded anti-apoptotic proteins and proteins homologous to BHRF1 expressed by herpes viruses and herpes-like viruses.

10 6. The method according to claim 1, wherein the monitoring step is by a method selected from the group consisting of co-precipitation, protein interactive trapping and ELISA.

1/6

GAGGATCTAC AGGGGACAAG TAAAGGCTAC ATCCAGATGC CGGGAATGCA CTGACGCCCA
60

TTCCTGGAAA CTGGGCTCCC ACTCAGCCCC TGGGAGCAGC AGCCGCCAGC CCCTCGGACC
120

TCCATCTCCA CCCTGCTGAG CCACCCGGGT TGGGCCAGGA TCCCGGCAGG CTGATCCCGT
180

CCTCCACTGA GACCTGAAAA ATG GCT TCG GGG CAA GGC CCA GGT CCT CCC
230

Met Ala Ser Gly Gln Gly Pro Gly Pro Pro
1 5 10

AGG CAG GAG TGC GGA GAG CCT GCC CTG CCC TCT GCT TCT GAG GAG CAG
278

Arg Gln Glu Cys Gly Glu Pro Ala Leu Pro Ser Ala Ser Glu Glu Gln
15 20 25

GTA GCC CAG GAC ACA GAG GAG GTT TTC CGC AGC TAC GTT TTT TAC CGC
326

Val Ala Gln Asp Thr Glu Glu Val Phe Arg Ser Tyr Val Phe Tyr Arg
30 35 40

CAT CAG CAG GAA CAG GAG GCT GAA GGG GTG GCT GCC CCT GCC GAC CCA
374

His Gln Glu Gln Glu Ala Glu Gly Val Ala Ala Pro Ala Asp Pro
45 50 55

GAG ATG GTC ACC TTA CCT CTG CAA CCT AGC AGC ACC ATG GGG CAG GTG
422

Glu Met Val Thr Leu Pro Leu Gln Pro Ser Ser Thr Met Gly Gln Val
60 65 70

GGA CGG CAG CTC GCC ATC ATC GGG GAC GAC ATC AAC CGA CGC TAT GAC
470

Gly Arg Gln Leu Ala Ile Ile Gly Asp Asp Ile Asn Arg Arg Tyr Asp
75 80 85 90

TCA GAG TTC CAG ACC ATG TTG CAG CAC CTG CAG CCC ACG GCA GAG AAT
518

Ser Glu Phe Gln Thr Met Leu Gln His Leu Gln Pro Thr Ala Glu Asn
95 100 105

GCC TAT GAG TAC TTC ACC AAG ATT GCC ACC AGC CTG TTT GAG AGT GGC
566

Ala Tyr Glu Tyr Phe Thr Lys Ile Ala Thr Ser Leu Phe Glu Ser Gly
110 115 120

Fig. 1

2/6

ATC AAT TGG GGC CGT GTG GTG GCT CTT CTG GGC TTC GGC TAC CGT CTG
 614
 Ile Asn Trp Gly Arg Val Val Ala Leu Leu Gly Phe Gly Tyr Arg Leu
 125 130 135

GCC CTA CAC GTC TAC CAG CAT GGC CTG ACT GGC TTC CTA GGC CAG GTG
 662
 Ala Leu His Val Tyr Gln His Gly Leu Thr Gly Phe Leu Gly Gln Val
 140 145 150

ACC CGC TTC GTG GTC GAC TTC ATG CTG CAT CAC TGC ATT GCC CGG TGG
 710
 Thr Arg Phe Val Val Asp Phe Met Leu His His Cys Ile Ala Arg Trp
 155 160 165 170

ATT GCA CAG AGG GGT GGC TGG GTG GCA GCC CTG AAC TTG GGC AAT GGT
 758
 Ile Ala Gln Arg Gly Gly Trp Val Ala Ala Leu Asn Leu Gly Asn Gly
 175 180 185

CCC ATC CTG AAC GTG CTG GTG GTT CTG GGT GTG GTT CTG TTG GGC CAG
 806
 Pro Ile Leu Asn Val Leu Val Val Leu Gly Val Val Leu Leu Gly Gln
 190 195 200

TTT GTG GTA CGA AGA TTC TTC AAA TCA TGA CTCCCAA GGGTGCCCTT
 853
 Phe Val Val Arg Arg Phe Phe Lys Ser
 205 210

TGGGTCCCGG TTCAGACCCC TGCCTGGACT TAAGCGAAGT CTTTGCCTTC TCTGTTCCTT
 913

TGCAGGGTCC CCCCTCAAGA GTACAGAAGC TTTAGCAAGT GTGCACTCCA GCTTCGGAGG
 973

CCCTGCGTGG GGGCCAGTCA GGCTGCAGAG GCACCTCAAC ATTGCATGGT GCTAGTGCCC
 1033

TCTCTCTGGG CCCAGGGCTG TGGCCGTCTC CTCCCTCAGC TCTCTGGGAC CTCCTTAGCC
 1093

CTGTCTGCTA GGCGCTGGGG AGACTGATAA CTTGGGGAGG CAAGAGACTG GGAGCCACTT
 1153

CTCCCCAGAA AGTGTTTAAC GGTTTTAGCT TTTTATAATA CCCTTGTGAG AGCCCATTC
 1213

CACCATTCTA CCTGAGGCCA GGACGTCTGG GGTGTGGGGA TTGGTGGGTC TATGTTCCCC
 1273

3/6

AGGATTCAGC TATTCTGGAA GATCAGCACC CTAAGAGATG GGA CTAGGAC CTGAGCCTGG
1333

TCCTGGCCGT CCCTAAGCAT GTGTCCCAGG AGCAGGACCT ACTAGGAGAG GGGGGCCAAG
1393

GTCCTGCTCA ACTCTACCCC TGCTCCCATT CCTCCCTCCG GCCATACTGC CTTTGCAGTT
1453

GGACTCTCAG GGATTCTGGG CTTGGGGTGT GGGGTGGGGT GGAGTCGCAG ACCAGAGCTG
1513

TCTGAACTCA CGTGTCAGAA GCCTCCAAGC CTGCCTCCCA AGGTCCTCTC AGTTCTCTCC
1573

CTTCCTCTCT CTTTATAGAC ACTTGCTCCC AACCCATTCA CTACAGGTGA AGGCTCTCAC
1633

CCATCCCTGG GGGCCTTGGG TGAGTGGCCT GCTAAGGCTC CTCCTTGCCC AGACTACAGG
1693

GCTTAGGACT TGGTTTGTTA TATCAGGGAA AAGGAGTAGG GAGTTCATCT GGAGGGTTCT
1753

AAGTGGGAGA AGGACTATCA ACACCACTAG GAATCCCAGA GGTGGATCCT CCCTCATGGC
1813

TCTGGCACAG TGTAATCCAG GGGTGTAGAT GGGGGAAGT TGAATACTG AACTCTGTTC
1873

CCCCACCCTC CATGCTCCTC ACCTGTCTAG GTCTCCTCAG GGTGGGGGGT GACAGTGCCT
1933

TCTCTATTGG CACAGCCTAG GGTCTTGGGG GTCAGGGGGG AGAAGTTCTT GATTGAGCCA
1993

AATGCAGGGA GGGGAGGCAG ATGGAGCCCA TAGGCCACCC CCTATCCTCT GAGTGTTTGG
2053

AAATAAACTG TGCAATCCCC TCAAAAAAAAA AACGGAGATC C
2094

4/6

10	20	30	40	50	60
TTT TAA TAT AAA TTA ATG TGC TCT ATT TAT AGA GAC AAT ACA TGA AAT ATA CTT AAT AAA					
AAA ATT ATA TTT AAT TAC ACG AGA TAA ATA TCT CTG TTA TGT ACT TTA TAT GAA TTA TTT					
70	80	90	100	110	120
AAT TCA AAT GTT ATA GAA CTG AAA AAG ATG AAA AGT AAA AAC AAC CTA TTC CCC AGA GGT					
TTA AGT TTA CAA TAT CTT GAC TTT TTC TAC TTT TCA TTT TTG TTG GAT AAG GGG TCT CCA					
130	140	150	160	170	180
AGC CAC TGT CCA TAG TTT CTA TTT TAG ATT CTT TCC TTT ATA CAA GAT TAT TAT AGC TTC					
TCG GTG ACA GGT ATC AAA GAT AAA ATC TAA GAA AGG AAA TAT GTT CTA ATA ATA TCG AAG					
190	200	210	220	230	240
TAT TTT TTG GTG TAT GAA CTG TAG TCC TAG AGG ATT TTA TTA GTT ATG AGT TCT ATA ACT					
ATA AAA AAC CAC ATA CTT GAC ATC AGG ATC TCC TAA AAT AAT CAA TAC TCA AGA TAT TGA					
250	260	270	280	290	300
AAG ATC CAT CAT CTT AGT TGC TAA GAA CGT AGA TAC TGA GAA CAT CAT TTA AAA AAA CAT					
TTC TAG GTA GTA GAA TCA ACG ATT CTT GCA TCT ATG ACT CTT GTA GTA AAT TTT TTT GTA					
310	320	330	340	350	360
TTT TGG CTG GCA CCT CAT GAT CAC TGG AGT CTC GCG GGT CCC TCA GGC TGC ACA GGG ACA					
AAA ACC GAC CGT GGA GTA CTA GTG ACC TCA GAG CGC CCA GGG AGT CCG ACG TGT CCC TGT					
370	380	390	400	410	420
AGT AAA GGC TAC ATC CAG ATG CTG GGA ATG CAC TGA CGC CCA TTC CTG GAA ACT GGG CTC					
TCA TTT CCG ATG TAG GTC TAC GAC CCT TAC GTG ACT GCG GGT AAG GAC CTT TGA CCC GAG					
430	440	450	460	470	480
CCA CTC AGC CCC TGG GAG CAG CAG CCG CCA GCC CCT CGG GAC CTC CAT CTC CAC CCT GCT					
GGT GAG TCG GGG ACC CTC GTC GTC GGC GGT CGG GGA GCC CTG GAG GTA GAG GTG GGA CGA					
490	500BamHI	510	520	530	540
GAG CCA CCC GGG TTG GGC CAG GAT CCC GGC AGG CTG ATC CCG TCC TCC ACT GAG ACC TGA					
CTC GGT GGG CCC AAC CCG GTC CTA GGG CCG TCC GAC TAG GGC AGG AGG TGA CTC TGG ACT					
550	560	570	580	590	600
AAA ATG GCT TCG GGG CAA GGC CCA GGT CCT CCC AGG CAG GAG TGC GGA GAG CCT GCC CTG					
TTT TAC CGA AGC CCC GTT CCG GGT CCA GGA GGG TCC GTC CTC ACG CCT CTC GGA CGG GAC					
M A S G Q G P G P P R Q E C G E P A L>					
610	620	630	640	650	660
CCC TCT GCT TCT GAG GAG CAG GTA GCC CAG GAC ACA GAG GAG GTT TTC CGC AGC TAC GTT					
GGG AGA CGA AGA CTC CTC GTC CAT CGG GTC CTG TGT CTC CTC CAA AAG GCG TCG ATG CAA					
P S A S E E Q V A Q D T E E V F R S Y V>					
670	680	690	700	710	720
TTT TAC CAC CAT CAG CAG GAA CAG GAG GCT GAA GGG GCG GCT GCC CCT GCC GAC CCA GAG					
AAA ATG GTG GTA GTC GTC CTT GTC CTC CGA CTT CCC CGC CGA CGG GGA CGG CTG GGT CTC					
F Y H H Q Q E Q E A E G A A A P A D P E>					

Fig. 2

5/6

```

      730          740          750 >NcoI      760          770          780
      *          *          *          *          *          *
ATG  GTC  ACC  TTA  CCT  CTG  CAA  CCT  AGC  AGC  ACC  ATG  GGG  CAG  GTG  GGA  CGG  CAG  CTC  GCC
TAC  CAG  TGG  AAT  GGA  GAC  GTT  GGA  TCG  TCG  TGG  TAC  CCC  GTC  CAC  CCT  GCC  GTC  GAG  CGG
M    V    T    L    P    L    Q    P    S    S    T    M    G    Q    V    G    R    Q    L    A>

      790          800          810          820          830          840
      *          *          *          *          *          *
ATC  ATT  GGG  GAC  GAC  ATC  AAC  CGA  CGC  TAT  GAC  TCA  GAG  TTC  CAG  ACC  ATG  TTG  CAG  CAC
TAG  TAA  CCC  CTG  CTG  TAG  TTG  GCT  GCG  ATA  CTG  AGT  CTC  AAG  GTC  TGG  TAC  AAC  GTC  GTG
I    I    G    D    D    I    N    R    R    Y    D    S    E    F    Q    T    M    L    Q    H>

>PstI  850          860          870          880          890          900
      *          *          *          *          *          *
CTG  CAG  CCC  ACG  GCA  GAG  AAT  GCC  TAT  GAG  TAC  TTC  ACC  AAG  ATT  GCC  TCC  AGC  CTG  TTT
GAC  GTC  GGG  TGC  CGT  CTC  TTA  CGG  ATA  CTC  ATG  AAG  TCG  TTC  TAA  CGG  AGG  TCG  GAC  AAA
L    Q    P    T    A    E    N    A    Y    E    Y    F    T    K    I    A    S    S    L    F>

      910          920          930          940          950          960
      *          *          *          *          *          *
GAG  AGT  GGC  ATC  AAT  TGG  GGC  CGT  GTG  GTG  GCT  CTT  CTG  GGC  TTC  AGC  TAC  CGT  CTG  GCC
CTC  TCA  CCG  TAG  TTA  ACC  CCG  GCA  CAC  CAC  CGA  GAA  GAC  CCG  AAG  TCG  ATG  GCA  GAC  CGG
E    S    G    I    N    W    G    R    V    V    A    L    L    G    F    S    Y    R    L    A>

      970          980          990          1000          1010          1020
      *          *          *          *          *          *
CTA  CAC  ATC  TAC  CAG  CGT  GGC  CTG  ACT  GGC  TTC  CTG  GGC  CAG  GTG  ACC  CGC  TTT  GTG  GTG
GAT  GTG  TAG  ATG  GTC  GCA  CCG  GAC  TGA  CCG  AAG  GAC  CCG  GTC  CAC  TGG  GCG  AAA  CAC  CAC
L    H    I    Y    Q    R    G    L    T    G    F    L    G    Q    V    T    R    F    V    V>

      1030          1040          1050          1060          1070          1080
      *          *          *          *          *          *
GAC  TTC  ATG  CTG  CAT  CAC  TGC  ATT  GCC  CGG  TGG  ATT  GCA  CAG  AGG  GGT  GGC  TGG  GTG  GCA
CTG  AAG  TAC  GAC  GTA  GTG  ACG  TAA  CGG  GCC  ACC  TAA  CGT  GTC  TCC  CCA  CCG  ACC  CAC  CGT
D    F    M    L    H    H    C    I    A    R    W    I    A    Q    R    G    G    W    V    A>

      1090          1100          1110          1120          1130          1140
      *          *          *          *          *          *
GCC  CTG  AAC  TTG  GGC  AAT  GGT  CCC  ATC  CTG  AAC  GTG  CTG  GTG  GTT  CTG  GGT  GTG  GTT  CTG
CGG  GAC  TTG  AAC  CCG  TTA  CCA  GGG  TAG  GAC  TTG  CAC  GAC  CAC  CAA  GAC  CCA  CAC  CAA  GAC
A    L    N    L    G    N    G    P    I    L    N    V    L    V    V    L    G    V    V    L>



      1150          1160          1170          1180          1190          1200
      *          *          *          *          *          *
TTG  GGC  CAG  TTT  GTG  GTA  CGA  AGA  TTC  TTC  AAA  TCA  TGA  CTC  CCA  AGG  GTG  CCT  TTG  GGG
AAC  CCG  GTC  AAA  CAC  CAT  GCT  TCT  AAG  AAG  TTT  AGT  ACT  GAG  GGT  TCC  CAC  GGA  AAC  CCC
L    G    Q    F    V    V    R    R    F    F    K    S    *>

      1210          1220          1230          1240          1250          1260
      *          *          *          *          *          *
TCC  CAG  TTC  AGA  CCC  CTG  CCT  GGA  CTT  AAG  CGA  AGT  CTT  TGC  CTT  CTC  TGC  TCC  TTG  CAG
AGG  GTC  AAG  TCT  GGG  GAC  GGA  CCT  GAA  TTC  GCT  TCA  GAA  ACG  GAA  GAG  ACG  AGG  AAC  GTC

      1270          1280 Hind3
      *          *
GT  CCC  CCC  TCA  AGA  GTA  CAG  AAG  CTT
CCA  GGG  GGG  AGT  TCT  CAT  GTC  TTC  GAA

```

6/6

**FIG. 3A****FIG. 3B**

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International Application No
PC./US 96/05639

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 G01N33/68 C07K14/47 C07K14/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 G01N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,95 05738 (MASSACHUSETTS INST TECHNOLOGY) 2 March 1995 Adenovirus E1B protein see page 10, line 1 - page 10, line 7 ---	1-6
A	CANCER RES. (1994), 54(10), 2808-11 CODEN: CNREA8; ISSN: 0008-5472, 1994, XP002007901 HICKISH, TAMAS ET AL: "Ultrastructural localization of BHRF1: an Epstein-Barr virus gene product which has homology with bcl-2" see the whole document --- -/--	1-6

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

9 July 1996

Date of mailing of the international search report

03. 10. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hoekstra, S

INTERNATIONAL SEARCH REPORT

International Application No

PC., US 96/05639

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VIROLOGY (1994), 201(2), 404-7 CODEN: VIRLAX;ISSN: 0042-6822, 1994, XP002007902 TARODI, BELA ET AL: "Epstein-Barr virus BHRF1 protein protects against cell death induced by DNA-damaging agents and heterologous viral infection" see the whole document ---	1-6
A	WO,A,95 05750 (UNIV WASHINGTON ;KORSMEYER STANLEY J (US)) 2 March 1995 see claim 8 ---	1-6
A	CELL, vol. 74, August 1993, page 609-619 XP002007903 OLTVAI, Z.N. ET AL.: "Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death" see the whole document ---	1-6
P,X	NATURE, vol. 374, no. 6524, 20 April 1995, pages 731-733, XP002007904 FARROW, S.N.: "cloning of a bcl-2 homologue by interaction with adenovirus E1B 19K" see page 733, left-hand column, last paragraph ---	1-6
P,A	NATURE, vol. 374, 20 April 1996, pages 736-739, XP002007905 MICHAEL J. KIEFER ET AL.: "Modulation of apoptosis by the widely distributed Bcl-2 homologue Bak" see page 739, right-hand column ---	1-6
P,A	WO,A,95 15084 (LXR BIOTECHNOLOGY INC ;KIEFER MICHAEL C (US); BARR PHILIP J (US)) 8 June 1995 cited in the application see page 16, line 25 - line 31 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PL./US 96/05639

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9505738	02-03-95	NONE	
WO-A-9505750	02-03-95	AU-B- 7604994 CA-A- 2170143	21-03-95 02-03-95
WO-A-9515084	08-06-95	AU-B- 1335195	19-06-95

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.